

Protocol of a systematic review with meta-analysis: Post-exercise hypotension effects of aquatic exercises on elderly

Tássia Magnabosco Sisoneto¹, Igor Moraes Mariano¹, Caroline Pereira Garcês¹, Ana Carolina Kanitz², Guilherme Morais Puga¹.

¹Research Group on Exercise, Women's Health and Cardiometabolic. Federal University of Uberlândia, Uberlândia-MG, 38400-678, Brazil.

²Multidisciplinary Research Group on Physical Exercise and Health. Federal University of Rio Grande do Sul, Porto Alegre-RS. 90010-150, Brazil.

**Corresponding author:* Faculty of Physical Education and Physiotherapy. Federal University of Uberlândia. Street Benjamin Constant, 1286. District: Nossa Senhora Aparecida. Uberlândia – MG, Brazil. CEP: 38400-678. Telefone/Fax: +55 34 3218-2901. Email: tassiamagnabosco@hotmail.com

TMS participated in the idealization, planning, original manuscript, and approval of the final version; IMM participated in the planning, review, and approval of the final version; CPG participated in the idealization, planning, review, and approval of the final version; ACK participated in the idealization, planning, review, and approval of the final version; GMP participated in the idealization, planning, review, and approval of the final version.

The authors declare that they have no conflicts of interest. Financial support: CAPES.

Abstract

Arterial hypertension (AH) is a chronic non-communicable disease of a multifactorial nature characterized by a consistent increase in blood pressure and is considered the main risk factor for the development of cardiovascular diseases. With aging, more than 70% of the elderly have AH. Aquatic exercise provides several benefits for middle-aged and elderly individuals, such as improving physical fitness and decreasing the heart rate of practitioners. However, the acute hypotensive responses of this type of exercise are not well documented in the literature, and in addition, the responses of blood pressure in the liquid environment during aquatic exercises vary according to body position, temperature, water height, and, mainly, with the intensity of the exercise. Thus, the objective of the scientific review is to answer the following question: What are the effects of aquatic exercises on acute blood pressure responses after performing these exercises in the elderly? For this, we will carry out a systematic review with meta-analysis in digital databases (EMBASE, PUBMED, Lilacs, SPORTDiscus, and Web of Science), of randomized and non-randomized clinical trials and cross-sectional studies carried out with elderly humans that indicate the values of post-exercise aquatic hypotension.

Keywords: Aquatic exercise, Blood Pressure, Hypotension, Elderly.

Background

Hypertension is a clinical condition associated with metabolic and hormonal disorders and target organ changes, characterized by increased blood pressure, and is the main risk factor for the development of cardiovascular diseases. In Brazil, more than 70% of the elderly population has AH, due to hemodynamic, hormonal, structural, and functional changes in the arterial vasculature. In addition, overweight and sedentary lifestyles can be mentioned (MALACHIAS et al., 2017; BARROSO et al., 2021).

It is well documented in the literature that aerobic, resistance and combined physical exercise can influence BP with just one session (CUNHA et al., 2013), causing the effect of post-exercise hypotension (PEH) that can generate a chronic reduction in BP over time (KEESE et al., 2011). Despite this, the hypotensive responses of aquatic exercise should be further investigated, since BP varies according to body position, temperature, and height of the water and, mainly, with the intensity of the exercise (ALBERTON et al., 2013).

Thus, the objective of the study is to answer the following question: What are the effects of aquatic exercises on acute blood pressure responses after performing these exercises in the elderly?

Methods

This systematic review with meta-analysis will follow the PRISMA guidelines and will be registered on the “PROSPERO” platform.

Eligibility criteria

Studies with the following categories will be eligible: 1) Population: Human, both sexes, elderly (>60 years); 2) Intervention: Acute session of any aquatic exercise modality; 3) Control: Immersion in water without exercise and resting on land; 4) Outcome of interest: values of blood pressure reduction after aquatic exercise (Systolic Blood Pressure, Diastolic Blood Pressure and/or Mean Blood Pressure); 5) Languages: English, Portuguese, and Spanish; 6) Study designs: randomized, non-randomized clinical trials and cross-sectional studies; 7) Publication dates: No time limit.

Excluded: literature reviews, meta-analysis, letters to the editor, observational studies, animal studies, studies in children, studies in adults, studies written in other languages, studies whose exercise intervention is performed on land, and studies that did not evaluate post-exercise hypotension.

Search strategy

Searches will be carried out in the following digital databases: EMBASE, PUBMED, Lilacs, SPORTDiscus, and Web of Science, in the references of the main articles and through manual search. If necessary, the authors of the studies will be contacted to request data.

The search will be divided into three categories of terms: Exercise, Post-exercise Hypotension, and Elderly. Within each category, terms will be separated by union operators (i.e. “OR”) and categories will be separated by parentheses and intersection operators (i.e. “AND”). All terms that will be added to the search are shown in table 1.

Table 1. Categorized search terms.

Exercise		Post-exercise hypotension	Elderly
Water exercise	Water running	Post-exercise hypotension	Aged
Water exercises	Running in water	Post exercise hypotension	Older adults
Exercises in water	Water walking	Hypotension	Elderly
Aquatic Exercise	Aquajogging	Blood pressure	Older
Aquatic Exercises	Shallow water walking	Systolic	Middle age
Water-based exercise	Aquatic environment	Diastolic	
Water-based exercises	Water resistance	Lowering blood pressure	
Water-based activities	Exercises in a pool	Acute blood pressure	
Water aerobics	Pool exercise	Mean blood pressure	
Water aerobic exercise	Water plyometric		
Water aerobic exercises	Swimming		
Deep water			

Study records

During the screening, eligibility, inclusion, and data extraction phases, studies will be evaluated in duplicate by 2 independent reviewers. After verifying the reviewers' responses, disagreements will be resolved by a third reviewer. These studies will be organized in the reference manager Rayyan (<https://www.rayyan.ai/>) and later recorded in a spreadsheet for data extraction and organization. If there are studies in which the data are represented only in graphs or figures without numerical representation, the data will be extracted by the software WebPlotDigitalizer (<https://automeris.io/WebPlotDigitizer/>).

Data extraction will include: 1) General description (identification code, author, year of publication, language, and study design); 2) Description of participants (gender, sample size, age); 3) Description of the exercise (modality, intensity, and volume); 4) Pool characteristics (water height and temperature); 5) Values referring to the primary outcome of the study (post-exercise hypotension in SBP, DBP, and MAP).

Risk of bias in individual studies

The risk of bias analysis will be performed using the tools of the Joanna Briggs Institute (<https://jbi.global/critical-appraisal-tools>). These will be presented in the results section in textual and/or graphical form and will help the explorer as texts from the studies in the discussion section. In addition, we will describe the conflicts of interest reported by the studies.

Data synthesis and quantitative approaches

Data analysis will be performed with the “R” programming language (R CORE TEAM, 2019) through the “meta” and “metafor” packages (VIECHTBAUER, 2010). They will be analyzed based on average differences. Kendall's tau and I^2 will be used as measures of heterogeneity. The meta-analysis values will be presented through “forest plots”. If the studies are sufficient, there will be a subgroup analysis based on the type of exercise intervention. Sensitivity analysis will be done through the search for outliers using the “externally standardized residuals” method (standard deviation values greater than 1.96 in the standardized residuals plot), and the search for influential points using the Difference of Fits methods (DFFITS) (identifying values above 1 or below -1), Covariance Ratio and Cook's Distance.

Publication bias analysis will be performed using a funnel plot and asymmetry hypothesis tests (Egger and Beggs).

References

ALBERTON, C.L. et al. Vertical ground reaction force during water exercises performed at different intensities. **Int J Sports Med.**, v. 34, p. 1-7, 2, 2013.

BARROSO, W. et al. Diretrizes Brasileiras de Hipertensão Arterial - 2020. **Arquivo Brasileiro de Cardiologia**, v. 116, n. 3, p. 516-658, 2021.

CUNHA, F.A. et al. Hipotensão pós-exercício induzida por treinamento aeróbio, de força e concorrente: aspectos metodológicos e mecanismos fisiológicos. **Rev HUPE**, v. 12, n. 4, 2013.

KEESE, F. et al. A comparison of the immediate effects of resistance, aerobic, and concurrent exercise on postexercise hypotension. **J Strength Cond Res**, v. 25, n. 5, 2011.

MALACHIAS, M. et al. 7ª Diretriz Brasileira de Hipertensão Arterial. **Brazilian Journal of Hypertension**, v. 24, n. 1, 2017.

R CORE TEAM. R: a language and environment for statistical computing [Internet]. **Vienna R Foundation for Statistical Computing**; 2019. Available from: <https://www.r-project.org>

VIECHTBAUER W. Conducting meta-analyses in R with the metafor package. **J Stat Softw.** v.36, p.1–48, 2010.